



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON THE SOLUTION OF EQUATIONS.

BY JOSEPH B. MOTT, ESQ., WORTHINGTON, MINNESOTA.

THE irrational roots of equations of all degrees may be determined by series. Thus, if

$$a = x + bx^2 + cx^3 + dx^4 + ex^5 + fx^6 + gx^7 + \&c. \quad (G)$$

we may find

$$\begin{aligned} x = a - ba^2 + (2b^2 - c)a^3 - (5b^3 - 5bc + d)a^4 + (14b^4 - 21b^2c + 6bd + 3c^2 - e)a^5 \\ - (42b^5 - 84b^3c + 28b^2d + 28bc^2 - 7be - 7cd + f)a^6 \\ + (132b^6 - 330b^4c + 120b^3d + 180b^2c^2 - 36b^2e - 72bcd + 8bf + 8ce - 12c^3 \\ + 4d^2 - g)a^7 - \&c., \quad (R) \end{aligned}$$

by assuming $x = a + Aa^2 + Ba^3 + Ca^4 + Da^5 + Ea^6 + Fa^7 + \&c.$, in (G).

Now let us take the following equations, viz.;

$$\begin{aligned} a &= x + 0x^2 + 0x^3 + 0x^4 + \&c., \text{ an eq'n of the 1st degree.} \\ a &= x + bx^2 + 0x^3 + 0x^4 + \text{ " " " " 2nd " } \\ a &= x + bx^2 + cx^3 + 0x^4 + \text{ " } \\ a &= x + 0x^2 + cx^3 + 0x^4 + \text{ " } \end{aligned} \left. \vphantom{\begin{aligned} a &= x + 0x^2 + 0x^3 + 0x^4 + \&c., \\ a &= x + bx^2 + 0x^3 + 0x^4 + \text{ " } \\ a &= x + bx^2 + cx^3 + 0x^4 + \text{ " } \\ a &= x + 0x^2 + cx^3 + 0x^4 + \text{ " } \end{aligned}} \right\} \text{eq'ns " " 3rd " } \\ \begin{aligned} a &= x + bx^2 + cx^3 + dx^4 + 0x^5 + \&c., \\ a &= x + 0x^2 + cx^3 + dx^4 + 0x^5 + \&c., \\ a &= x + 0x^2 + 0x^3 + dx^4 + 0x^5 + \&c., \end{aligned} \left. \vphantom{\begin{aligned} a &= x + bx^2 + cx^3 + dx^4 + 0x^5 + \&c., \\ a &= x + 0x^2 + cx^3 + dx^4 + 0x^5 + \&c., \\ a &= x + 0x^2 + 0x^3 + dx^4 + 0x^5 + \&c., \end{aligned}} \right\} \text{eq's of 4th degree.} \\ \begin{aligned} a &= x + bx^2 + cx^3 + dx^4 + ex^5 + 0x^6 + \&c., \\ a &= x + bx^2 + cx^3 + 0x^4 + ex^5 + 0x^6 + \&c., \\ a &= x + 0x^2 + cx^3 + 0x^4 + ex^5 + 0x^6 + \&c., \\ a &= x + 0x^2 + 0x^3 + 0x^4 + ex^5 + 0x^6 + \&c., \end{aligned} \left. \vphantom{\begin{aligned} a &= x + bx^2 + cx^3 + dx^4 + ex^5 + 0x^6 + \&c., \\ a &= x + bx^2 + cx^3 + 0x^4 + ex^5 + 0x^6 + \&c., \\ a &= x + 0x^2 + cx^3 + 0x^4 + ex^5 + 0x^6 + \&c., \\ a &= x + 0x^2 + 0x^3 + 0x^4 + ex^5 + 0x^6 + \&c., \end{aligned}} \right\} \text{eq'ns of 5th deg., \&c.}$$

From which it appears that equation (G) is a complete expression for an equation of any degree, and that its series may be made to vanish at any term, so as to give any degree, by giving zero values to such of the coefficients $b, c, d, \&c.$ as the case may require.

The series in (R), however, will not end after the first degree, but its terms may be greatly contracted, in many cases, by cancelling all of those having zero coefficients. Thus, if in both (G) and (R) we make $c=0, d=0, e=0, \&c.$, we get,

$$a = x + bx^2, \quad (G')$$

$$x = a - ba^2 + 2b^2a^3 - 5b^3a^4 + 14b^4a^5 - 42b^5a^6 + 132a^7 - \&c. \quad (R')$$

1. Given $x^2 + 10x = 1$, or $\frac{1}{10} = x + \frac{1}{10}x^2$. Put $a = \frac{1}{10}$ and $b = \frac{1}{10}$ in (R'), then $x = \frac{1}{10} - \frac{1}{10^3} + \frac{2}{10^5} - \frac{5}{10^7} + \&c. = .0990195$

2. $x^2 + 10x = 5$, or $\frac{1}{2} = x + \frac{1}{10}x^2$. Here $a = \frac{1}{2}, b = \frac{1}{10}$; therefore

$$x = \frac{1}{2} - \frac{1}{10 \cdot 2^2} + \frac{2}{10^3 \cdot 2^3} - \frac{5}{10^5 \cdot 2^4} + \frac{14}{10^7 \cdot 2^5} - \frac{42}{10^9 \cdot 2^6} + \frac{132}{10^{11} \cdot 2^7} = .477225$$

3. $15x^2 - 50x + 7 = 0$, or $\frac{7}{50} = x - \frac{5}{10}x^2$. Here $a = \frac{7}{50}, b = -\frac{5}{10}$;

$$\therefore x = \frac{7}{50} + \frac{3 \cdot 7^2}{10 \cdot 50^2} + \frac{2 \cdot 3^2 \cdot 7^3}{10^2 \cdot 50^3} + \frac{5 \cdot 3^3 \cdot 7^4}{10^3 \cdot 50^4} + \&c.$$

4. Given $x^2+x=1$. Here $a=1$ and $b=1$; $\therefore x=1-2+5-14$, &c.

5. $x^2+x=100$. $a=100$, $b=1$; $x=100-2.100^2+5.100^3$ —&c.

From Examples 1, 2 and 3, we see that the values of a and b in formula (R') when applied should be small fractions or the series will not converge rapidly, but may diverge as in Ex's 4 and 5. Hence direct application of the formula would only be practically useful in comparatively few cases.

There is, however, an indirect method, as presented in my Math. Key, which makes the formula applicable to all cases. Thus, in the equation $x^2+x=100$, we may soon find by trial that x lies between 9 and 10, then put $x=\frac{9}{2}+y$, and we have $(\frac{9}{2}+y)^2+(\frac{9}{2}+y)=100$, or $\frac{1}{4.20}=y+\frac{1}{20}y^2$;

$$\therefore x=\frac{19}{2}+y=\frac{19}{2}+\frac{1}{4.20}-\frac{1}{4^2.20^3}+\frac{2}{4^3.20^5}$$
—&c. = 9.51249.

6. Given, $x^2+6x=8$, to find x . As the value is near 1, let $x=1+y$. Then the eq. may be written $\frac{1}{8}=y+\frac{1}{8}y^2$, and $x=1+y=1+\frac{1}{8}-\frac{1}{8^3}+\frac{2}{8^5}$, &c.

7. Required the square root of 2, or the value of x in the eq'n $x^2=2$.

Let $x=1.4+y$, as 1.4 is near the value of x . Then $(1.4+y)^2=1.96+2.8y+y^2=2$, or $\frac{1}{70}=y+\frac{5}{14}y^2$. Developing y by (R') as before, we find

$$x=\frac{14}{10}+\frac{1}{70}-\frac{5}{14.70^3}+\frac{2.5^2}{14^2.70^5}$$
—&c.

In all the examples here given other series might be found that would converge much faster, by taking a nearer value of x before transforming.

In the solution of cubic equations, we may consider the values of all the letters zero, except a , b , c and x , in both (G) and (R), and by then cancel'g all zero terms we have $a=x+bx^2+cx^3$; and for its root,

$x=a-ba^2+(2b^2-c)a^3-(5b^3-5bc)a^4+(14b^4-21b^2c+3c^2)a^5$ —&c.; (R'')
or, when $b=0$, $a=x+cx^3$, and

$$x=a-ca^3+3c^2a^5-12c^3a^7$$
—&c. (R''')

8. Given $x^3-3x^2+10x=1$, or $\frac{1}{10}=x-\frac{3}{10}x^2+\frac{1}{10}x^3$. Let $a=\frac{1}{10}$, $b=\frac{3}{10}$ and $c=\frac{1}{10}$ in (R'); then $x=\frac{1}{10}+\frac{3}{10^3}+\frac{8}{10^5}-\frac{15}{10^7}$ —&c. = .1031, nearly.

9. Given, $x^4-3x^2+75x=10000$, to find one root of the equation.

As upon trial x is found nearly equal 10, let $x=9.9+y$, and the equation becomes, $-.0139676=y+.15y^2$ +&c. Developing by (R') we get

$$x=9.9+y=9.9-.0139676-.0000293-.0000001$$
—&c. = 9.8860027.

When the fractions of the new equation have large terms it is best to reduce to decimals, as in the last case.

10. Given, $x^5-30x^4+340x^3-1800x^2+4384x=3841$, to find all the r'ts.

As the several values of x are near to 2, 4, 6, 8 and 10, respectively; \therefore substitute successively $x_1=2+y$, $x_2=4+y$, $x_3=6+y$, $x_4=8+y$ and $x_5=10+y$. The new equations will then be

